


GR ASSIGNMENTS 01



1. METRICS, LINE ELEMENTS AND COORDINATE TRANSFORMATIONS

Under a coordinate transformation from Cartesian (or inertial) coordinates $\xi^a = \xi^a(x^\mu)$, the Euclidean metric δ_{ab} or the Minkowski metric η_{ab} transforms to a new metric $g_{\mu\nu}$ in such a way that proper distances are invariant,

$$(\delta_{ab} \text{ or } \eta_{ab})d\xi^a d\xi^b = g_{\mu\nu}dx^\mu dx^\nu \quad \Leftrightarrow \quad g_{\mu\nu} = J_\mu^a J_\nu^b (\delta_{ab} \text{ or } \eta_{ab}) \quad , \quad (1)$$

with $J_\mu^a = \partial\xi^a/\partial x^\mu$ the Jacobi matrix of the transformation.

(a) 2-DIMENSIONAL METRICS

Consider the 2-dimensional line element

$$ds^2 = (dy^1)^2 + (dy^2)^2 + 2ady^1 dy^2 \quad , \quad (2)$$

with $a \in \mathbb{R}$ a real constant parameter.

- Show that this metric is non-degenerate for $a \neq \pm 1$.
- Show that for $a^2 < 1$ the metric is related to the standard Euclidean metric $ds^2 = (dx^1)^2 + (dx^2)^2$ by the coordinate transformation

$$x^1 = \sqrt{1 - a^2}y^1 \quad , \quad x^2 = ay^1 + y^2 \quad . \quad (3)$$

- Show that for $a^2 > 1$ the metric is related to the standard Lorentzian metric $ds^2 = -(dt)^2 + (dx)^2$ by a coordinate transformation.

(b) RINDLER METRIC

Consider the (1+1)-dimensional Minkowski metric with line element $ds^2 = -dt^2 + dx^2$, and the *Rindler coordinates* (T, X) , defined by

$$t = X \sinh T \quad , \quad x = X \cosh T \quad . \quad (4)$$

- Determine the metric or line element in these coordinates.
- Show that the lines of constant T or constant X are straight lines through the origin or hyperbolae (in a (t, x) -diagram) respectively.

2. GEODESICS I

(a) GEODESICS AND EULER-LAGRANGE EQUATIONS:

Show that the Euler-Lagrange equations

$$\frac{d}{d\tau} \frac{\partial \mathcal{L}}{\partial \dot{x}^\mu} - \frac{\partial \mathcal{L}}{\partial x^\mu} = 0 \quad , \quad (5)$$

for the Lagrangian

$$\mathcal{L} = \frac{1}{2} g_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} \quad (6)$$

are the geodesic equations

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\nu\lambda} \frac{dx^\nu}{d\tau} \frac{dx^\lambda}{d\tau} = 0 \quad , \quad (7)$$

where the *Christoffel symbols* $\Gamma^\mu_{\nu\lambda}$ associated to a metric $g_{\mu\nu}$ are defined by

$$\Gamma^\mu_{\nu\lambda} = g^{\mu\rho} \Gamma_{\rho\nu\lambda} = \frac{1}{2} g^{\mu\rho} (g_{\rho\nu,\lambda} + g_{\rho\lambda,\nu} - g_{\nu\lambda,\rho}) \quad (8)$$

(b) \mathcal{L} IS A CONSTANT OF MOTION:

Show that \mathcal{L} is constant along any geodesic, i.e. that

$$\frac{d}{d\tau} \left(g_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} \right) = 0 \quad (9)$$

for $x^\mu(\tau)$ a solution of the geodesic equation.

[As an aside: what is the symmetry that, by Noether's theorem, gives rise to this constant of motion?]