


## GR ASSIGNMENTS 06



### 1. RIEMANN CURVATURE TENSOR AND DIFFERENTIAL IDENTITIES

In the course, we defined the Riemann curvature tensor via the commutator of covariant derivatives,

$$[\nabla_\alpha, \nabla_\beta]V^\gamma = R^\gamma_{\delta\alpha\beta}V^\delta \quad , \quad [\nabla_\alpha, \nabla_\beta]T^{\gamma\delta} = R^\gamma_{\epsilon\alpha\beta}T^{\epsilon\delta} + R^\delta_{\epsilon\alpha\beta}T^{\gamma\epsilon} \quad , \quad (1)$$

etc. and we defined its contractions, the Ricci tensor  $R_{\alpha\beta} = R^\gamma_{\alpha\gamma\beta}$  and the Ricci scalar  $R = g^{\alpha\beta}R_{\alpha\beta}$ . The Riemann curvature tensor has the symmetries

$$R_{\alpha\beta\gamma\delta} = -R_{\alpha\beta\delta\gamma} = -R_{\beta\alpha\gamma\delta} \quad , \quad R_{\alpha[\beta\gamma\delta]} = 0 \Leftrightarrow R_{\alpha\beta\gamma\delta} + R_{\alpha\gamma\delta\beta} + R_{\alpha\delta\beta\gamma} = 0 \quad (2)$$

These symmetries also imply that the Ricci tensor is symmetric,  $R_{\alpha\beta} = R_{\beta\alpha}$ . [see section 8.3 of the lecture notes for proofs and make sure that you understand the details!]

- (a) Like any linear operator, the covariant derivative  $\nabla_\alpha$  satisfies the Jacobi identity

$$[\nabla_\alpha, [\nabla_\beta, \nabla_\gamma]] + \text{cyclic permutations} = 0 \quad . \quad (3)$$

Show that this implies the *Bianchi identity*

$$\nabla_\alpha R_{\mu\nu\beta\gamma} + \text{cyclic permutations in } (\alpha, \beta, \gamma) = 0 \quad (4)$$

- (b) By double-contraction of the Bianchi identity, deduce the *contracted Bianchi identity*

$$\nabla_\alpha(2R^\alpha_\gamma - \delta^\alpha_\gamma R) = 0 \quad (5)$$

and show that this is equivalent to the statement that the *Einstein tensor*  $G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R$  has vanishing covariant divergence,  $\nabla^\alpha G_{\alpha\beta} = 0$ .

- (c) By double-contracting the expression for  $[\nabla_\alpha, \nabla_\beta]T^{\gamma\delta}$ , show that for any tensor  $T^{\alpha\beta}$  one has

$$[\nabla_\alpha, \nabla_\beta]T^{\alpha\beta} = 0 \quad . \quad (6)$$

- (d) Use this result to show that the Maxwell equations  $\nabla_\alpha F^{\alpha\beta} = -J^\beta$  imply that the current is covariantly conserved,

$$\nabla_\alpha F^{\alpha\beta} = -J^\beta \quad \Rightarrow \quad \nabla_\beta J^\beta = 0 \quad . \quad (7)$$

**Remark:** In all these equations indices are lowered and raised with the metric and its inverse:  $R_{\alpha\beta\gamma\delta} = g_{\alpha\lambda}R^\lambda_{\beta\gamma\delta}$ ,  $\nabla^\alpha = g^{\alpha\rho}\nabla_\rho$ ,  $R^\alpha_\gamma = g^{\alpha\beta}R_{\beta\gamma}$  etc.

## 2. CURVATURE OF A CLASS OF 2-DIMENSIONAL METRICS

Consider the 2-dimensional line element

$$ds^2 = dx^2 + f(x)^2 d\phi^2 . \quad (8)$$

Show that the Ricci tensor and Ricci scalar curvature of this metric are

$$R_{\alpha\beta} = -(f''/f)g_{\alpha\beta} \quad , \quad R(x) = -2f''(x)/f(x) . \quad (9)$$

**Hint:** Calculate the Christoffel symbols from the Euler-Lagrange equations. Then determine the one independent component of the Riemann tensor, e.g.  $R^x_{\phi x \phi}$ , and then deduce the Ricci tensor and Ricci scalar from this.

**Remark:** In particular, for the Euclidean metric and the standard metrics on the sphere and the hyperboloid one finds

$$f(x) = \begin{cases} x & (R^2) \\ \sin x & (S^2) \\ \sinh x & (H^2) \end{cases} \quad \Rightarrow \quad R = \begin{cases} 0 \\ +2 \\ -2 \end{cases} \quad (10)$$

In 2 dimensions,  $R$  is related to the *Gauss Curvature*  $K$  of a surface by  $K = R/2$  so that  $K = 0, \pm 1$  in these examples.