

## KFT SOLUTIONS 02

### 1. ACTION FOR A FREE PARTICLE

The action is

$$S[x] = -mc^2 \int d\tau = -mc^2 \int d\lambda (d\tau/d\lambda) \equiv \int d\lambda L_\lambda \quad (1)$$

with  $L_\lambda = -mc^2(d\tau/d\lambda) = -mc(-\eta_{\alpha\beta}x'^\alpha x'^\beta)^{1/2}$ .

(a) The momentum is

$$\begin{aligned} p_\alpha &= \frac{\partial L_\lambda}{\partial x'^\alpha} = -mc \frac{1}{2} \left( -\eta_{\alpha\beta} x'^\alpha x'^\beta \right)^{-1/2} \left( -2\eta_{\alpha\beta} x'^\beta \right) \\ &= mc \eta_{\alpha\beta} \left( \frac{cd\tau}{d\lambda} \right)^{-1} \frac{dx^\beta}{d\lambda} = m\eta_{\alpha\beta} \frac{dx^\beta}{d\tau} = mu_\alpha \end{aligned} \quad (2)$$

or  $p^\alpha = mu^\alpha$ . In an inertial system with coordinates  $(x^0 = ct, x^k)$ , and with  $v^k = dx^k/dt$  one has

$$p^0 = m\gamma(v)c = E/c \quad , \quad p^k = m\gamma(v)v^k \quad . \quad (3)$$

(b) The Euler-Lagrange equation is

$$\frac{d}{d\lambda} \frac{\partial L_\lambda}{\partial x'^\alpha} - \frac{\partial L_\lambda}{\partial x^\alpha} = \frac{d}{d\lambda} \frac{\partial L_\lambda}{\partial x'^\alpha} = 0 \quad (4)$$

and

$$\frac{d}{d\lambda} \frac{\partial L_\lambda}{\partial x'^\alpha} = \frac{d\tau}{d\lambda} \frac{d}{d\tau} mu_\alpha = \left( m\eta_{\alpha\beta} \frac{d\tau}{d\lambda} \right) \frac{d^2 x^\beta}{d\tau^2} = 0 \quad \Leftrightarrow \quad \frac{d^2 x^\beta}{d\tau^2} = 0 \quad . \quad (5)$$

(c) The Lagrangian is

$$L_t = -mc^2 \frac{d\tau}{dt} = -mc^2 \sqrt{1 - \vec{v}^2/c^2} = -mc^2 \gamma(v)^{-1} \quad . \quad (6)$$

Thus the canonical momenta are

$$p_k^{(c)} = \frac{\partial L_t}{\partial v^k} = (-mc^2)\gamma(v)(-v^k/c^2) = m\gamma(v)v^k = p^k \quad , \quad (7)$$

and the canonical Hamiltonian is

$$\begin{aligned} H &= p_k^{(c)} v^k - L_t = m\gamma(v)\vec{v}^2 + mc^2\gamma(v)^{-1} \\ &= m\gamma(v)(\vec{v}^2 + c^2(1 - \vec{v}^2/c^2)) = m\gamma(v)c^2 = E = cp^0 \quad . \end{aligned} \quad (8)$$